



I'M NO ANGEL



FIND A WAY OUT OF YOUR GOUT HELL BY FIRST ASKING YOUR DOCTOR THE RIGHT QUESTIONS.

It's important to talk openly with your doctor. Knowing what information and questions to bring up can help you have a productive conversation about your gout experience and treatment options.



TELL YOUR DOCTOR

◆ In the last 12 months I have experienced gout flares:

- 0–1 times
- 2–3 times
- 4–7 times
- 8+ times

◆ When I have a flare, I usually manage the pain by taking:

- OTC pain medication
- Colchicine
- Corticosteroids
- Other: _____

ASK YOUR DOCTOR

◆ What were the results of my last two uric acid level tests?

- Date: _____ Level: _____
- Date: _____ Level: _____

◆ Did I reach the American College of Rheumatology recommended uric acid level of <6 mg/dL?

- Yes
- No

KEEP YOUR GOAL IN MIND

Avoiding certain foods is a must, but diet alone may not be enough. The American College of Rheumatology guidelines recommend a goal serum uric acid (sUA) of less than 6 mg/dL for all patients with gout. If your sUA level is greater than 6 mg/dL, your gout is uncontrolled.



DUZALLO is used to lower high uric acid levels in the blood in people with gout when allopurinol alone has not worked well enough. It works differently because it **reduces** uric acid production and **helps eliminate** uric acid from the body.

Before taking DUZALLO, tell your healthcare provider about all of your medical conditions.

Important Safety Information

DUZALLO can cause serious side effects, including:

Kidney Problems: Some people taking DUZALLO may have kidney problems, such as a sudden decrease in kidney function (acute kidney failure). Your healthcare provider may do blood tests to check your kidneys while you are taking DUZALLO.

IF YOUR DOCTOR THINKS DUZALLO® IS APPROPRIATE FOR YOU, EXPLORE SAVINGS ON YOUR PRESCRIPTION



You may be eligible to receive a **30-day supply** for a **\$0 co-pay**.* After the first 30 days, you could receive a 30-day refill for as little as **\$15** or a 90-day refill for as little as **\$30**.*

Visit [DUZALLO.com/Savings](https://www.duzallo.com/Savings) to check your eligibility.

Offer not valid for patients enrolled in government healthcare programs.

*Maximum savings limit applies; patient out-of-pocket expense may vary. Offer not valid for patients enrolled in Medicare, Medicaid, or other federal or state healthcare programs. This offer is not valid for cash-paying patients. Please see Program Terms, Conditions, and Eligibility Criteria at [DUZALLO.com/Savings](https://www.duzallo.com/Savings) for details.

Important Safety Information (cont.)

Serious Skin Rash and Serious Allergic Reactions:

Severe and potentially life-threatening allergic reactions and skin reactions, including Stevens-Johnson syndrome, toxic epidermal necrolysis, and exfoliative dermatitis, have been reported in some people taking allopurinol, one of the components of DUZALLO. **Stop taking DUZALLO and get emergency medical help if you have any of the following symptoms:**

- skin rash; blistering of the skin, mouth, nose, and genitals
- painful red or purple skin that looks burned and peels off
- joint pain; swollen glands
- red, painful, watery eyes; persistent itching; hives
- fever

Do not take DUZALLO if you have severe kidney problems, have received a kidney transplant or are on dialysis, have a fast breakdown of cancer cells that can lead to high uric acid (tumor lysis syndrome), have a rare inherited condition that causes too much uric acid in the blood (Lesch-Nyhan syndrome), or had a skin rash or other allergic reaction after taking allopurinol.

Liver Problems: Liver problems can happen in people who take DUZALLO. Call your healthcare provider if you have any of the following symptoms: loss of appetite, weight loss, feeling very tired, nausea or vomiting, yellowing of your skin or the white part of your eyes, itching, dark or brown (tea-colored) urine, or pain on the upper right side of your stomach.

Heart Problems: People who take DUZALLO can have serious heart problems, including heart attack and stroke. It is not known that DUZALLO causes these problems.

Blood Problems: DUZALLO can affect your bone marrow and cause low red blood cells, white blood cells, and platelets. If your blood cell counts become very low, you can get infections or have bleeding problems. Taking DUZALLO with certain other medicines can increase your chance of having blood problems.

Drowsiness: DUZALLO can cause you to be drowsy. Do not drive, operate heavy machinery, or do other dangerous activities until you know how DUZALLO affects you.

The most common side effects of DUZALLO include headache, flu, higher levels of blood creatinine (a measure of kidney function), heartburn (acid reflux), skin rash, nausea, and diarrhea. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all of the possible side effects of DUZALLO. For more information, ask your healthcare provider or pharmacist.

What is DUZALLO?

DUZALLO is a prescription medicine that contains 2 medicines, a URAT1 inhibitor called lesinurad and a xanthine oxidase inhibitor called allopurinol. DUZALLO is used to lower uric acid levels in the blood in people with gout, when allopurinol alone has not worked well enough.

Please see [Medication Guide](#) and full [Prescribing Information](#), including [Boxed Warning](#).

Ironwood® and the three-leaf design are registered trademarks of Ironwood Pharmaceuticals, Inc.

DUZALLO® and its logo are registered trademarks of AstraZeneca or its affiliates and are used herein under license.

©Ironwood Pharmaceuticals, Inc. 2018
All rights reserved. PP-DUZ-US-00202 03/18

